

Forest Service

Southern Forest Experiment Station


New Orleans, Louisiana


Resource Bulletin SO-151



## Distribution Maps for Midsouth Tree Species

Roy C. Beltz, and Daniel F. Bertelson





## Distribution Maps for Midsouth Tree Species

Roy C. Beltz and Daniel F. Bertelson

The Midsouth is an important timber-producing region, with a wide variety of sites and species. In addition to timber production, increasing demands for non-timber amenities are placed on the region's forests. These maps indicate the distribution of individual species recorded in surveys of the Midsouth conducted by the U.S. Department of Agriculture, Forest Servee. Included are important timber species plus several species that are important primarily from a wildlife, aesthetic, or other nontimber perspective.

Approximately 17,500 forest plots are remeasured during a complete survey cycle of the seven Midsouth states (Alabama, Arkansas, Louisiana, Mississippi, Oklahoma, Tennessee, and Texas). These plots, each about 1 acre in size and representing roughly 9 square miles, are depicted in figure 1 in their approximate geographic location.


In the late 1940's and early 1950's, survey plots were established throughout the Midsouth using a 3-mile grid spacing. Permanent plots were installed at forested grid intersections. In some instances, the grid was overlaid on an entire survey unit; in others, single counties (parishes in Louisiana) were overlaid. Coordinates used to depict plot locations were developed by referencing one central point in each county. The latitude and longitude of all plots in the county were subsequently developed from the central coordinate on the basis of the 3- by 3-mile spacing. Thus, the coordinates are crude and serve only to position the plot attributes relative to one another within the Midsouth. Moreover, the actual ground location of a plot may differ from a precise 3- by 3-spacing because of establishment procedures and changes in conditions over several decades.

Prominent features of the region are immediately apparent from the distribution of forest plots. Sparsely forested areas are apparent in the Missisppi Delta in Arkansas, Mississippi, and Louisiana, primarily due to the predominance of agricultural land use. Similarly, the black prairie region extending from southeast Alabama westward and northward to northeast Mississippi is sparsely forested. The same is true of the Red and Arkansas River bottoms.


Other areas are sparsely forested for varying physical reasons. The coastal marshes of Louisiana and Texas, for example, do not support forests. The western extremity of southern forests is limited by the average annual rainfail amounts in Texas and Oklahoma. Dots represent plots where the forest is capable of growing 20 or more cubic feet of wood per acre per year.

To be recorded in the survey, a species must be capable of growing to fit the definition of a tree; i.e., 15 feet tall and having a central stem. Species that do not occur in the region, or that were not identified by survey field crews, were not considered. Some species were grouped because of the difficulty in identifying them or because they occur infrequently in the survey.

Several sources of error could affect the species distribution maps. Since surveys are done year-round, and certain species are hard to identify without leaves, some species identification errors are bound to occur. Seedlings are especially subject to this type of error. Errors are also made when trying to distinguish minor differences between very similar species. In addition, some species are known to exist in the Midsouth, but, unless they happen to be growing on one of the 3- by 3-mile sample plots, they will not be represented in the mapped data.




 $\label{prop:continuous} Figure \ 1. -Location \ of \ forest \ inventory \ and \ analysis \ plots \ on \ timber land.$ 




## **Softwoods**


Southern redcedar (Juniperus silicicola)


Eastern redcedar (Juniperus virginiana)







Shortleaf pine (Pinus echinata)















Eastern white pine (Pinus strobus)






Loblolly pine (Pinus taeda)




Virginia pine (Pinus virginiana)







Eastern hemlock (Tsuga canadensis)



## Hardwoods


Florida maple (Acer barbatum)







Boxelder (Acer negundo)





























Bumelia, chittamwood (Bumelia spp.)





























Eastern redbud (Cercis canadensis)







Common persimmon (Diospyros virginiana)




American beech (Fagus grandifolia)




White ash (Fraxinus americana)




Green ash (Fraxinus pennsylvanica)






Kentucky coffeetree (Gymnocladus dioicus)










Butternut (Juglans cinerea)










Sweetgum (Liquidambar styraciflua)










Osage-orange (Maclura pomifera)










Bigleaf magnolia (Magnolia macrophylla)





Sweetbay (Magnolia virginiana)















Swamp tupelo (Nyssa sylvatica var. biflora)








**Sourwood** (Oxydendrum arboreum)





















Swamp white oak (Quercus bicolor)















Bluejack oak (Quercus incana)







Laurel oak (Quercus laurifolia)












Swamp chestnut oak (Quercus michauxii)


Chinkapin oak (Quercus muehlenbergii)








Water oak (Quercus nigra)















Chestnut oak (Quercus prinus)









Black oak (Quercus velutina)




Live oak (Quercus virginiana)





Sassafras (Sassafras albidum)







American basswood (Tilia americana)






Cedar elm (Ulmus crassifolia)





Slippery elm (Ulmus rubra)



Sparkleberry (Vaccinium arboreum)

# **Distribution Map Index**

Page

#### Softwoods

#### Scientific Name

### Juniperus silicicola ......3 Pinus elliotti ......4 Pinus glabra .....4 Pinus palustris ......5 Pinus pungens ......5 Pinus strobus ......6 Common Name Cypress ......8 Eastern hemlock ......8 Longleaf pine ......5 Slash pine .....4 Spruce pine ......4 Table Mountain pine ......5

#### Hardwoods

#### Scientific Name

	Page
Acer barbatum	9
Acer negundo	9
Acer nigrum	
Acer rubrum	
Acer saccharinum	
Acer saccharum	
Aesculus spp.	
Ailanthus altissima	
Betula alleghaniensis	. 13
Betula lenta	
Betula nigra	
Bumelia spp.	.14
Caroinus caroliniana	
Carya aquatica	
Carya illinoensis	.16
Carya spp	
Castanea dentata	
Castanea spp.	
Catalpa spp.	18
Celtis laevigata	18
Celtis occidentalis	
Cercis canadensis	
Cornus florida	
Cornus portua Crataegus spp.	
Diospyros virgininana	21
Pagus grandifolia	
Fraxinus americana	22
Fraxinus americana Fraxinus pennsylvanica	22
Fraunus pennsylvanica	22
Gleditsia spp	
Gymnocladus dioicus	. 23
Rex opaca	
Juglans cinerea	
Juglans nigra	
Liquidambar styraciflua	
Liriodendron tulipifera	
Maciura pomifera	
Magnolia acuminata	
Magnolia grandiflora	
Magnolia macrophylla	
Magnolia virginiana	
Malus spp	
Melia azedarach	. 29
Morus spp.	
Nyssa aquatica	
Nyssa sylvatica	
Nyssa sylvatica vat. biflora	.31
Ostrya virginiana	.32
Oxydendrum arboreum	32

#### Hardwoods—Continued

#### Scientific Name—Continued Common Name—Continued Page Page Blackiack oak ...... 42 Quercus bicolor ......36 Ouercus faicata ......38 Boxelder ..... 9 Bur oak ...... 41 Quercus laurifolia .....40 Quercus lyrata ......41 Quercus macrocarpa ......41 Cedar elm .....52 Quercus michauxii ......42 Quercus nigra ..... 43 Chinkapin ...... 17 Cottonwood ......34 Ouercus rubra ......46 Quercus shumardii .....46 Durand oak ...... 37 Quercus virginiana ......48 Robinia pseudoacacia ......49 Hawthorn ...... 20 Sassafras albidum ......50 Hickory ...... 16 Ulmus alata ......51 Ulmus americana ......51 Ulmus crassifolia ......52 Ulmus rubra ......52 Vaccinum arboreum ......53 Common Name Nuttall oak ...... 44 American basswood ......50 Overcup oak ......41 American chestnut 17 American holly .......24 American sycamore ......34

## Hardwoods-Continued

## Common Name-Continued

## Common Name—Continued

	Page
Sassafras	50
Scarlet oak	37
Shingle cak	39
Shumard oak	46
Silver maple	11
Slippery elm	52
Sourwood	32
Southern magnolia	27
Southern red oak	38
Sparkleberry	53
Sugarberry	18
Sugar maple	11
Swamp chestnut oak	42
Swamp tupelo ,	31
Swamp white oak	36
F	70

	Page
Sweet birch	. 13
Sweetgum	. 25
Tree-of-heaven	. 12
Turkey cak	.40
Water-elm	.33
Water hickory	. 15
Water oak	. 43
Water tupelo	.30
White ash	
White oak	.36
Willow	. 49
Willow oak	. 45
Winged elm	.51
Yellow-poplar	
Yellow birch	

Beltz, Roy C.; Bertelson, Daniel F. 1990. Distribution maps for Midsouth tree species. Resour. Bull. SO-151.New Orleans, L.A: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 56 p.

Depicts the location of tree species recorded in the U.S. Department of Agriculture, Forest Service, forest survey of the Midsouth,

Keywords: Maps, species, tree.

Persons of any race, color, national origin, sex, age, religion, or with any handicapping condition are welcome to use and enjoy all facilities, programs, and services of the USDA. Discrimination in any form is strictly against a gency policy, and should be reported to the Secretary of Agriculture, Washington, DC 20250.